

UPR-0525T

September, 1992

PARTICLE WIDTH EFFECT ON CP -VIOLATING ASYMMETRIES

JIANG LIU

Department of Physics, University of Pennsylvania, Philadelphia, PA 19104

ABSTRACT

CP -violating asymmetries calculated from the Breit-Wigner approximation for unstable particle propagators violate CPT . A formalism satisfying CPT invariance and unitarity is presented. Applications are given to t decays. For the decay $t \rightarrow b\nu_\tau\bar{\tau}$ in a class of CP -violating Higgs models the CP -violating asymmetry turns out to be 10^{-4} times smaller than the previously reported results.

PACS#: 11.30.Er, 13.20.Jf, 13.30.Ce.

Final-state interactions play an important role in the determination of CP and T violation. A test for CP violation is to compare the partial decay rates of a particle and its antiparticle. In this case final-state interactions are necessary since in their absence the partial decay rates are equal from CPT invariance even if CP is violated.¹

Examples of interest are those involving an intermediate unstable particle, in which the final-state interaction enters into the standard calculation as the width of the unstable particle. In this case the Breit-Wigner approximation is usually introduced, under which the W gauge boson propagator is given in the unitary gauge by

$$iD_{\mu\nu}(k)_{BW} = \frac{-i}{k^2 - M_W^2 + i\Gamma_W M_W} \left[g_{\mu\nu} - \frac{k_\mu k_\nu}{M_W^2} \right], \quad (1)$$

where M_W and Γ_W are the W mass and total width, respectively. The subscript BW refers to the Breit-Wigner approximation.

This approximation violates CPT invariance and its effect can be very large in calculating delicate quantities such as CP -violating asymmetries in weak decays.

Consider the difference of the following partial rates of the t decays

$$\Delta_{b\nu_\tau\bar{\tau}} \equiv \Gamma(\bar{t} \rightarrow \bar{b}\bar{\nu}_\tau\tau) - \Gamma(t \rightarrow b\nu_\tau\bar{\tau}) \quad (2)$$

in the CP -violating Higgs model of Weinberg.² We consider this example because (1) there is a considerable interest in these decays,^{3,4} and (2) in this case the CPT violation effect due to the Breit-Wigner approximation is most drastic.

The interaction between the charged-Higgs bosons and the fermions of the

model is given by⁵

$$L(H_{1,2}^+) = \frac{g}{\sqrt{2}M_W} \sum_{i=1}^2 H_i^+ \left[\bar{U} \left(Y_i M_u K L + X_i K M_d R \right) D - Z_i \bar{\nu} M_\ell R \ell \right], \quad (3)$$

where M_u , M_d and M_ℓ are the mass matrices for the up quarks, down quarks, and leptons, respectively. K is the Kobayashi-Maskawa matrix,⁶ L and R are the helicity projection operators. X_i , Y_i and Z_i are dimensionless parameters arising from the diagonalization of the mass matrix of the charged scalars. They depend on three mixing angles and one CP -violating phase.

Neglecting the contributions from the neutral Higgs sector,⁷ the lowest order CP -violating asymmetry $\Delta_{b\nu_\tau\bar{\tau}}$ is produced by the interference of the tree-level W and charged-Higgs exchange diagrams. Assuming, for simplicity, that the contributions from H_2^+ are negligible and that the final-state interaction effect due to the charged-Higgs boson width is insignificant, a straightforward calculation shows that the Breit-Wigner approximation result is

$$\Delta_{b\nu_\tau\bar{\tau}} = \frac{-g^4}{512\pi^3} \left(\frac{m_\tau}{M_W} \right)^2 \frac{\tilde{\Gamma}_W}{M_W} \frac{m_t^3 M_W^2 \text{Im}(Y_1 Z_1)}{(M_W^2 - M_{H_1^+}^2)^2 + \Gamma_W^2 M_W^2} F\left(\frac{M_W^2}{m_t^2}, \frac{M_{H_1^+}^2}{m_t^2}\right), \quad (4)$$

where

$$F(x, y) = \left[\frac{y}{x} - 1 \right] \left[(1-x)^2 \ln \frac{1-x}{x} - (1-y)^2 \ln \frac{1-y}{y} + (x-y) \right], \quad (5)$$

and

$$\tilde{\Gamma}_W = \Gamma_W - \Gamma(W \rightarrow \nu_\tau \bar{\tau}) \approx \frac{8}{9} \Gamma_W. \quad (6)$$

In reaching Eq. (4) we have ignored the b -quark mass and assumed $\Gamma_W M_W \ll |M_W^2 - M_{H_1^+}^2|$. The dependence of $\Delta_{b\nu_\tau\bar{\tau}}$ on $\tilde{\Gamma}_W$ arises because the same final state

rescattering does not contribute to its CP -violating asymmetry.⁸ An important feature of Eq. (4) is that $\Delta_{b\nu_\tau\bar{\tau}}$ is quadratic in m_τ . One power of m_τ comes from $L(H_1^+)$ (see Eq. (3)). The other is the price which one pays for the interference of a vector-current (due to W^+) and a scalar-current (due to H_1^+).

Under the Breit-Wigner approximation $\Delta_{b\nu_\tau\bar{\tau}}$ is the largest CP asymmetry that can be produced by the standard weak currents and the interaction given by Eq. (3). The analogous asymmetries of other t decays are completely negligible: the other semileptonic decays are suppressed by either m_μ^2 or m_e^2 , where $m_{\mu(e)}$ is the muon(electron) mass; the lowest order hadronic decays conserve CP if we ignore the small down-type quark masses. It then follows that there would be no other CP -violating asymmetries big enough to cancel $\Delta_{b\nu_\tau\bar{\tau}}$, and thus the lifetime of t would not be the same as the lifetime of $\bar{t}$, in violation of CPT invariance.

CPT violation occurred in the calculation discussed above is a consequence of the Breit-Wigner approximation, which misses an important final-state interaction phase.

Consider the W propagator in the unitary gauge. The tree-level result is

$$iD_{\mu\nu}^{(0)}(k) = \frac{-i}{k^2 - M_W^2} \left[g_{\mu\nu} - \frac{k_\mu k_\nu}{M_W^2} \right]. \quad (7)$$

Parameterizing the one-loop W self-energy as

$$\Pi_{\mu\nu}(k) = g_{\mu\nu}F_1(k^2) + (k^2g_{\mu\nu} - k_\mu k_\nu)F_2(k^2), \quad (8)$$

we find that the regularized (but unrenormalized) W propagator is

$$\begin{aligned} iD_{\mu\nu}^{(1)}(k) &= -i \left[(k^2g_{\mu\nu} - k_\mu k_\nu) - M_W^2g_{\mu\nu} - \Pi_{\mu\nu}(k^2) \right]^{-1} \\ &= \frac{-i[1 - F_2(k^2)]^{-1}}{k^2 - [M_W^2 + F_1(k^2)][1 - F_2(k^2)]^{-1}} \left[g_{\mu\nu} - k_\mu k_\nu \frac{1 - F_2(k^2)}{M_W^2 + F_1(k^2)} \right]. \end{aligned} \quad (9)$$

In analogous to the optical theorem, the unitarity of S -matrix implies a relation between $Im\Pi_{\mu\nu}\epsilon^\mu\epsilon^\nu|_{k^2=M_W^2}$ and Γ_W :

$$ImF_1(M_W^2) + M_W^2 ImF_2(M_W^2) = -\Gamma_W M_W, \quad (10)$$

where ϵ_μ is the W polarization vector. It then follows from the standard on-shell renormalization procedure that

$$iD_{\mu\nu}^{(1)}(k) = \frac{-iQ(k^2)}{k^2 - M_W^2 + i\Gamma_W M_W - \Sigma_R(k^2)} \left[g_{\mu\nu} - \frac{k_\mu k_\nu}{M_W^2 - i\Gamma_W M_W + \Sigma_R(k^2)} \right], \quad (11)$$

where the mass renormalization correction is $\Delta M_W^2 = ReF_1(M_W^2) + M_W^2 ReF_2(M_W^2)$ and

$$\begin{aligned} Q(k^2) &= [1 - F_2(k^2) + F_2(M_W^2)]^{-1}, \\ \Sigma_R(k^2) &= [F_1(k^2) - F_1(M_W^2)] + M_W^2 [F_2(k^2) - F_2(M_W^2)] \end{aligned} \quad (12)$$

are the renormalized weak charge correction and self-energy, respectively. Complications in the longitudinal part of the propagator are usually neglected in the literature, for in most of the practical cases the $k_\mu k_\nu$ term either vanishes (if it couples to a conserved current) or is very small (if it couples to fermions with small masses).

Compared to the Breit-Wigner approximation (Eq. (1)), the result given by Eq. (11) has an important difference: the width term $i\Gamma_W M_W$ is always in company with the mass term M_W^2 everywhere in the formula, not just in the denominator of the propagator. Other modifications include (1) an over all screening factor $Q(k^2)$ for the weak charge, and (2) a renormalized self-energy $\Sigma_R(k^2)$. Both $Q(k^2)$ and $\Sigma_R(k^2)$ can have a final-state interaction phase, but their effects are likely to be

less significant as they vanish at the pole (see below). The position of the pole of the propagator remains the same, i.e.,

$$k^2 = M_W^2 - i\Gamma_W M_W, \quad (13)$$

and the relations $\Sigma_R(k^2)|_{k^2=M_W^2-i\Gamma_W M_W} = 0$ and $Q(k^2)|_{k^2=M_W^2-i\Gamma_W M_W} = 1$ hold to the order of interest to us in the perturbation expansion.⁹

Turn to the calculation of the CP -violating asymmetries in the t decays. For the case at hand, effects arising from the weak charge screening factor $Q(k^2)$ and the self-energy $\Sigma_R(k^2)$ are negligible, and hence to a good approximation the W propagator can be written as

$$iD_{\mu\nu}(k) \approx \frac{-i}{k^2 - M_W^2 + i\Gamma_W M_W} \left[g_{\mu\nu} - \frac{k_\mu k_\nu}{M_W^2 - i\Gamma_W M_W} \right]. \quad (14)$$

The phase in the longitudinal part of the propagator is crucial in restoring CPT invariance of the calculation. It introduces a correction to $\Delta_{b\nu_\tau\bar{\tau}}$ given by Eq. (4). The correction has the same amplitude as Eq.(4) but with the opposite sign. As a consequence, the CPT -violating result (Eq. (4)) cancels completely, as it should.

It is not totally surprising that the final-state interaction phase in the $k_\mu k_\nu$ term of the W propagator has such a big effect. It has been noticed by the previous studies (Refs. 3, 4) that under the Breit-Wigner approximation the interference term of the squared amplitudes vanishes for an on-shell W . This does not happen to the correction terms however, even though they appear to be suppressed by a factor Γ_W/M_W .

Because of the large cancellation imposed by CPT invariance, the size of the CP -violating asymmetries induced by the mechanism discussed above turns out to

be purely academic. The order-of-magnitude of the leading terms is approximately

$$\begin{aligned} \Delta_{b\nu_\tau\bar{\tau}} &= -\Delta_{bc\bar{s}} \\ &\sim \frac{-g^4}{512\pi^3} \left(\frac{m_\tau}{M_W}\right)^2 \left(\frac{m_c}{M_W}\right)^2 \frac{\tilde{\Gamma}_W}{M_W} \frac{m_t^3 M_W^2 |Y_1|^2 \text{Im}(Y_1 Z_1)}{(M_W^2 - M_{H_1^+}^2)^2 + \Gamma_W^2 M_W^2} f\left(\frac{M_W^2}{m_t^2}, \frac{M_{H_1^+}^2}{m_t^2}, \frac{m_\tau^2}{m_t^2}, \frac{m_c^2}{m_t^2}\right), \end{aligned} \quad (15)$$

where $\Delta_{bc\bar{s}} = \Gamma(\bar{t} \rightarrow \bar{b}\bar{c}s) - \Gamma(t \rightarrow bc\bar{s})$, m_c is the c -quark mass, and f is a function of order unity or less and symmetric under the exchange of m_τ and m_c . Although the calculations of the decay rates $\Gamma(t \rightarrow b\nu_\tau\bar{\tau})$ and $\Gamma(t \rightarrow bc\bar{s})$ are very different, there is a complete correspondence imposed by CPT invariance when we calculate the rate difference. Compared to Eq. (4), the unitarized and CPT invariant result (Eq. (15)) has an additional suppression factor $\sim |Y_1|^2 m_c^2 / M_W^2$ and is about 10^{-4} times smaller.

In general, neglecting the phase in the $k_\mu k_\nu$ part of the W propagator will have an important implication for CP asymmetries generated from interactions involving an interference that requires a small mass insertion. The effect is expected to be small for interactions¹⁰ involving currents of the same type, unless the final-state interaction phase factors.

The propagator of an unstable fermion with mass m and width Γ is

$$iS(k) = \frac{i}{\not{k} - m + i\Gamma/2 - \Sigma_R(k^2)} \approx i \frac{\not{k} + m - i\Gamma/2}{k^2 - m^2 + i\Gamma m}, \quad (16)$$

where $\Sigma_R(k^2)$ is the renormalized self-energy of the fermion. The approximation in the second step of Eq. (16) corresponds to neglecting $\Sigma_R(k^2)$. The phase factor $i\Gamma/2$ in the numerator has so far been ignored in the literature.¹¹ Its consequence should be examined carefully. Little change is expected from an unstable scalar, whose propagator is the same as the standard Breit-Wigner form if one ignores its renormalized self-energy.

In conclusion, we have shown that the standard Breit-Wigner approximation for an unstable particle violates *CPT*. The effect can be large in calculating delicate quantities such as *CP*-violating asymmetries in some theoretical models. Formalisms satisfying *CPT* invariance and unitarity are derived. For the *CP*-asymmetry discussed in more detail in this paper, their applications show that the result is about 10^{-4} times smaller than that obtained from the Breit-Wigner approximation.

Acknowledgements:

I wish to thank M. Cvetič, B. Kayser, P. Langacker and L. Wolfenstein for valuable discussions and suggestions. I would also like to thank Aspen Center for Physics for hospitality during the initial stage of this work. This research was supported in part by the U. S. Department of Energy under the contract DE-ACO2-76-ERO-3071 and the SSC National Fellowship.

REFERENCES

1. For a review see: L. Wolfenstein, *Ann. Rev. Nucl. Part. Sci.* 36, 137 (1986).
2. S. Weinberg, *Phys. Rev. Lett.* 37, 657 (1976).
3. G. Eilam, R. R. Mendel, R. Migeron, and A. Soni, Technion-Israel Inst. Tech. report, TECHNION-PH-92-10, (1992).
4. R. Cruz, B Grzdkowski, and J. F. Gunion, UC Davis report, UCD-92-15, (1992).
5. C. H. Albright, J. Smith, and S.-H. Tye, *Phys. Rev.* D21, 711 (1980).
6. M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* 49, 652 (1973).
7. Interactions involving a charged- and a neutral-Higgs boson can provide a dominant contribution to the CP -violating asymmetry provided $m_t \geq m_b + M_{H^+}$, where H^+ is either H_1^+ or H_2^+ or both. This possibility will not be discussed in this paper.
8. L. Wolfenstein, *Phys. Rev.* D43, 151 (1991).
9. The effect of particle width on the definition of particle mass becomes slightly more complicated in higher orders of perturbation expansion: A. Borrelli, M. Consoli, L. Maiani and R. Sisto, *Nucl. Phys.* B333, 357 (1990); A. Borrelli, L. Maiani and R. Sisto, *Phys. Lett.* B244, 117 (1990); A. Sirlin, *Phys. Rev. Lett.* 67, 2127 (1991); R. Stuart, *Phys. Lett.* B262, 113 (1991); G. Valencia and S. Willenbrook, *Phys. Rev.* D40, 853 (1990); *Phys. Lett.* B259, 373 (1991).
10. G. Eilam, J. L. Hewett, and A. Soni, *Phys. Rev. Lett.* 67, 1979 (1991); *ibid* 68, 2103 (1992); J. M. Soares, *Phys. Rev. Lett.* 68, 2102 (1992).

11. M. Nowakowski and A. Pilaftsis, Mod. Phys. Lett. A4, 821 (1989); Z. Phys. C42, 449 (1989); Mod. Phys. Lett. 6A, 1933 (1991); A. Pilaftsis, Z. Phys. C47, 95 (1990).